Tree Topping Hurts!

Topping is severely cutting limbs larger than 3" in diameter to stubs within the tree's crown so as to remove the normal canopy and disfigure the tree.

Reasons not to top:

■ **Starvation** Good pruning practices rarely remove more than 1/2 to 1/3 of the crown, which in turn does not seriously interfere with the ability of a tree's leafy crown to manufacture food. Topping removes so much of the crown that it upsets an older tree's well-developed crown-to-root ratio and temporarily cuts off its food-making ability.

■ **Shock** A tree's crown is like an umbrella that shields much of the tree from the direct rays of the sun. By suddenly removing this protection, the remaining bark tissue is so exposed that scalding may result. There may also be a dramatic effect on neighboring trees and shrubs. If these thrive in shade and the shade is removed, poor health or death may result.

■ **Ugliness** A topped tree is a disfigured tree. Even with its regrowth it never regains the grace and character of its species. The landscape and the community are robbed of a valuable asset.

■ Rapid new growth The goal of topping is usually to control the height and spread of a tree. Actually, it has just the opposite effect. The resulting sprouts (often called water sprouts) are far more numerous than normal new growth and they elongate so rapidly that the tree returns to its original height in a very short time — and with a far denser crown.

Insects and disease The large stubs of a topped tree have a difficult time forming callus. The terminal location of these cuts, as well as their large diameter, prevent the tree's chemically based natural defense system from

doing its job. The stubs are highly vulnerable to insect invasion and the spores of decay fungi. If decay is already present in the limb, opening the limb will speed the spread of the disease.

a tree is much easier than applying the skill and judgement of good pruning. Therefore, topping may cost less in the short run. However, the true costs of topping are hidden. These include: reduced property value, the expense of removal and replacement if the tree dies, the loss of other trees and shrubs if they succumb to changed light conditions, the risk of liability from weakened branches, and increased future maintenance.

■ **Weak limbs** At best, the wood of a new limb that sprouts after a larger limb is cut is more weakly attached than a limb that develops more normally. If rot exists or develops at the severed end of the limb, the weight of the sprout makes a bad situation even worse.

■ **Tree death** Some older trees are more tolerant to topping than others. Beeches, for example, do not sprout readily after severe pruning and the reduced foliage most surely will lead to death of the tree.



Alternatives to topping

- Start out right by planting trees that will fit your available space when they reach maturity
- Prune properly and regularly. A light pruning every three years will keep your tree in healthy condition.
- Crown reduction, making the tree canopy smaller using proper pruning techniques, can be performed by a certified arborist.

For more information, contact

North Carolina Urban Forest Council PO Box 25852, Winston-Salem, NC 27114-5852 919-733-2162 x248 www.NCUFC.org



Funding for this project was provided in part through an Urban and Community Forestry Grant from the North Carolina Division of Forest Resources, Department of Environment and Natural Resources, in cooperation with the USDA Forest Service, Southern Region.